

H.S. Johnson

Rhodora

JOURNAL OF THE
NEW ENGLAND BOTANICAL CLUB

Conducted and published for the Club, by

BENJAMIN LINCOLN ROBINSON, Editor-in-Chief

MERRITT LYNDON FERNALD
HOLLIS WEBSTER
CARROLL WILLIAM DODGE } Associate Editors

WILLIAM PENN RICH, Publication Committee

Vol. 30.

December, 1928.

No. 360.

CONTENTS:

Notes on Panicum in Rhode Island. <i>J. F. Collins</i>	229
The Identification of Linnaean Species. <i>K. K. MacKenzie</i>	232
Trisetum in America (concluded). <i>Father Louis-Marie</i>	237
Errata.....	246
Index.....	247

Boston, Mass.

300 Massachusetts Ave.



Providence, R. I.

Preston and Rounds Co.

RHODORA.—A monthly journal of botany, devoted primarily to the flora of New England. Price, \$2.00 per year, postpaid (domestic and foreign); single copies (if available) 20 cents. Volumes 1-8 or single numbers from them can be supplied at somewhat advanced prices which will be furnished on application. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be gladly received and published to the extent that the limited space of the journal permits. Forms will be closed five weeks in advance of publication. Authors (of more than one page of print) will receive 25 copies of the issue in which their contributions appear. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to

B. L. ROBINSON, 3 Clement Circle, Cambridge, Mass.

Subscriptions, advertisements, and business communications to

W. P. RICH, 300 Massachusetts Avenue, Boston, Mass.

Entered at Boston, Mass., Post Office as Second Class Mail Matter.

BOTANICAL BOOKS,

New and Second Hand,

PRESTON & ROUNDS CO.,

Providence, R. I.

CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS, 1885 TO DATE.

For American taxonomists and all students of American plants the most important supplement to the Index Kewensis, this catalogue in several ways exceeds the latter work in detail, since it lists not only the flowering plants, but ferns and other vascular cryptogams, and includes not merely genera and species, but likewise subspecies, varieties and forms. A work of reference invaluable for larger herbaria, leading libraries, academies of sciences, and other centers of botanical activity. Issued quarterly, at \$22.50 per 1000 cards.

GRAY HERBARIUM of Harvard University,

Cambridge, Mass., U. S. A.

CHECK LIST OF GRAY'S MANUAL, 7th EDITION, compiled by M. A. DAY. Leatherette. Pocket size. Invaluable for collector's memoranda and herbarium records. Published and sold by the GRAY HERBARIUM, Cambridge, Mass. Price postpaid 20 cts. each. Ten copies \$1.50.

MEMOIRS OF THE GRAY HERBARIUM. A series of illustrated quarto papers issued at irregular intervals, sold separately.

Vol. II. Persistence of Plants in unglaciated Areas of Boreal America, by M. L. Fernald, 102 pages. Aug. 1925. \$2.00

Gray Herbarium of Harvard University, Cambridge, Mass.

Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. by 3-4 in. 1 year \$4.00, 6 months \$2.50.

Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

Vol. 30.

December, 1928.

No. 360.

NOTES ON CERTAIN SPECIES OF PANICUM OCCURRING IN OR NEAR RHODE ISLAND

J. FRANKLIN COLLINS

IN February, 1928, Mr. C. A. Weatherby examined the specimens of *Panicum* in my herbarium, and also others which were unnamed and unmounted. He was kind enough to revise the names of a few of the species and identify all of the unnamed ones, except in the case of a few immature specimens. Somewhat later I became interested in checking up the available published records of distribution of the different species, particularly records of their occurrence in Rhode Island and adjacent Massachusetts and Connecticut. This was followed by a check-up of specimens in the herbarium of the New England Botanical Club and the Gray Herbarium. The results were unexpectedly interesting from the point of view of the presence or absence of certain species in Rhode Island. These results are here briefly summarized as it is thought probable that they may be of interest to other readers of RHODORA, especially to those who collect in southern New England. The lists of species enumerated below include more particularly those in the herbaria mentioned which have been reported or collected recently in Rhode Island. They also include such as are known to occur in adjacent states and might reasonably be expected in Rhode Island. The asterisk (*) indicates species that are recorded in manuals in a general way as occurring in Rhode Island, but are not specifically included from the state in Hitchcock & Chase, "North American Species of *Panicum*" (1910). Specimens in the Gray Herbarium are indicated by (G), those in the herbarium of the New England Botanical Club by (NE), and those in the herbarium of the writer by (C).

(1) Reported from Connecticut and Long Island but not from Rhode Island. *P. amarum* Ell., *P. Boscii* var. *molle* (Vasey) H. & C.

(2) Reported from Long Island, Connecticut and Massachusetts but not from Rhode Island. *P. Addisonii* Nash., *P. Commonsianum* Ashe, *P. lucidum* Ashe.

(3) Reported from Long Island and Massachusetts but not Rhode Island. *P. Wrightianum* Scribn. (*P. minutulum* Desv.).

(4) Reported from Connecticut and Massachusetts but not Rhode Island. *P. Bicknellii* Nash., **P. Boscii* Poir., **P. xanthophysum* Gray.

(5) Reported from Connecticut but not Rhode Island. *P. pseudo-pubescent* Nash, *P. stipitatum* Nash.

(6) Reported from Massachusetts but not Rhode Island. *P. calliophyllum* Ashe, *P. commutatum* Schultes, *P. Clutei* Nash.

The following notes refer primarily to specimens seen in one or more of the herbaria mentioned above.

P. ALBEMARLENSE Ashe. First reported from Rhode Island by Fernald (RHOD. 24: 98) in 1922. It is now known from the towns of Warwick, Portsmouth (Prudence Island), Westerly, New Shoreham (Block Island), and Little Compton (NE)—all towns bordering on salt water.

P. ANNULUM Ashe. Reported from New Jersey southward and westward. It has been collected by Fernald in Harwich, Mass. (NE. G).

P. AUBURNE Ashe. Reported from Virginia southward. It has been collected on Cape Cod, Mass. (NE. G).

P. BARBULATUM Mx. Reported from Mass. and Conn. It has been collected in Johnston, Warwick (NE), and Westerly, R. I. (C).

**P. CAPILLARE* L. Collected in Warwick and Westerly, R. I. (NE).

P. CAPILLARE var. *occidentale* Rydb. (*P. barbipulvinatum* Nash). Reported from Illinois westward. Probably naturalized eastward (RHOD. 21: 110). Reported by Bicknell (RHOD. 16: 82) in 1914 from Marthas Vineyard, Mass., and Long Island, N. Y. Since then specimens have been identified as this variety from Prince Edward Island, New Brunswick, and Nova Scotia, and practically across the country to the Pacific coast (G). In Rhode Island it has been collected in East Providence (Williams, 1903) and Block Island (Fernald, Long and Torrey, 1913) (G).

**P. COLUMBIANUM* Scribn. Collected in Warwick, Little Compton (C), and Charlestown, R. I. (NE).

P. COLUMBIANUM var. *thinium* H. & C. Reported from Massachusetts, New Jersey, and southward. It has been collected in Connecticut (NE), and in Hopkinton, R. I. (C).

P. DEPAUPERATUM var. *PSILOPHYLLUM* Fernald. Described as a *n. var.* in 1921 (RHOD. 23: 193). It has been collected in Johnston, Smithfield, Glocester (C), Providence, Warwick (NE. G), Cumberland, East Providence, Coventry, Hopkinton, Westerly, and Middletown, R. I. (NE).

**P. DICHOTOMIFLORUM* Mx. Collected in Providence (NE. C), Warwick, Westerly, and Block Island, R. I. (NE).

P. DICHOTOMIFLORUM var. *PURITANORUM* Svenson. Described as a *n. var.* in 1920 (RHOD. 22: 154) from Cape Cod, Mass. Collected in South Kingstown, R. I., in 1914 by Collins and Fernald (NE).

**P. DICHOTOMUM* L. Reported from Warwick, R. I., in 1908 (RHOD. 12: 216). It is widely distributed in Rhode Island, e. g., in Providence, Lincoln, Johnston (C), Hopkinton (NE. C), Warwick, South Kingstown, Westerly (NE).

P. LANUGINOSUM Ell. Reported from New Jersey southward. It was collected on Block Island, R. I., in 1913 by Fernald, Long and Torrey (NE).

**P. LINEARIFOLIUM* Scribn. Collected in Cumberland in 1917 by Knowlton, and in Foster, R. I., in 1922 by Eaton and Fassett (NE).

**P. LINEARIFOLIUM* var. *WERNERI* (Scribn.) Fernald. Collected in Glocester (NE), Wickford (G), and Block Island, R. I. (NE. G).

P. LONGIFOLIUM Torr. Reported by H. & C. (p. 106) from "Shannock, R. I." This village is situated partly in Richmond and partly in Charlestown. In addition to this station it is known from Hopkinton (NE. C), Westerly and Richmond, R. I. (NE).

P. MATTAMUSKEETENSE Ashe. Reported from Long Island southward. Erroneously reported from Massachusetts in 1901 (RHOD. 3: 114 and 15: 57). It has been collected in Massachusetts (NE. G) and by the writer in South Kingstown, R. I., in 1927 (NE. G. C).

**P. MILIACEUM* L. To be expected on any dumping ground. Collected in Little Compton, East Providence (C), and Providence, R. I. (NE. C).

P. OLIGOSANTHES Schultes. Reported from New Jersey to Texas, and in Massachusetts in 1913 (RHOD. 15: 58 and 64). Collected on Block Island, R. I., in 1913 by Fernald and Long (NE. G).

P. PHILADELPHICUM Bernh., as redefined by Fernald (RHOD. 21: 112). Reported from Connecticut. Collected in Hopkinton, R. I., in 1919 by Ware, Woodward and Harger (NE).

P. SCOPARIOIDES Ashe. Reported from Connecticut (H. & C., p. 239, and RHOD. 15: 66). It has been collected at Sharon, Mass. (NE).

P. SCOPARIUM Lam. Reported from Cape Cod (H. & C., p. 295) and Marthas Vineyard (RHOD. 16: 82), Mass., and from New Jersey southward. Collected on Block Island, R. I., in 1916 by Collins, Gravatt and Spaulding (NE).

P. SUBVILLOSUM Ashe. Reported from Massachusetts and Con-

necticut. Collected by Fernald, Long and Torrey on Block Island, R. I., in 1913 (G).

**P. VILLOSISSIMUM* Nash. Collected in Hopkinton, R. I., in 1919 by Fernald, Woodward and Collins (NE).

BROWN UNIVERSITY.

CONCERNING THE PROPER IDENTIFICATION OF LINNAEAN SPECIES, ESPECIALLY THOSE BASED ON MATERIAL COLLECTED BY CLAYTON

KENNETH K. MACKENZIE

ONE of the great basic collections of American plants is the collection made by Clayton in Virginia. This is preserved in the British Museum. Of it Gray (Scientific Papers of Asa Gray 2: 9-10) says: "But still more important is the herbarium of Clayton, from whose notes and specimens Gronovius edited the 'Flora Virginica.' Many Linnaean species are founded on the plants here described for which this herbarium is alone authentic; for Linnaeus, as we have already remarked, possessed very few of Clayton's plants. The collection is nearly complete, but the specimens were not well prepared, and are not therefore always in perfect preservation." "From Gronovius, Linnaeus had received a very small number of Clayton's plants, previous to the publication of the 'Species Plantarum'; but most of the species of the 'Flora Virginica' were adopted or referred to other plants on the authority of the descriptions alone." (l. c. 6.)

We must also bear in mind that Linnaeus had actively assisted Gronovius in the *Flora Virginica*, which was published in 1739-1743. "Other work of Linnaeus in Leyden consisted * * *. He also helped Gronovius with his 'Flora Virginica' in which Linnaeus's principles were embodied." (Jackson, *Linnaeus* 165; *Pulteney, Linnaeus* 49.) Gronovius in the preface to his work (p. 3) acknowledged the assistance of Linnaeus as follows:

"Nullus igitur dubitavi specimina plantarum cum perspicacissimo Linnaeo examinare; utinam reliqua etiam cum doctissimo viro ad examen revocare mihi licuisset."

So when we find Linnaeus in 1753 in his *Species Plantarum* constantly referring to Gronovius' *Flora Virginica*, these references are to a work in which he had assisted and to a collection with which he was personally thoroughly familiar. And these references are of

the most definite nature, because they are to the specific specimens with which Gronovius and Linnaeus worked.

The situation with respect to the Linnaean herbarium on the other hand is very unsatisfactory and very exasperating. Its condition is fully discussed by the late Dr. B. D. Jackson (Proc. Linnean Society, 1912 Appendix). Information as to collector, place and time of receipt of a specimen is very often absent, and only arbitrary signs or such abbreviations as "K" for Kalm usually appear. Sometimes there are no data at all. The name and the number in the first edition of the *Species Plantarum* usually are given, additions made afterwards being given letters.

From three lists (not entirely accurate apparently) which are preserved we know what species Linnaeus had represented in his herbarium in 1753, in 1755 and in 1767. However, we cannot be sure that the specimens he had in those years are the specimens existing now, for he was constantly adding to his herbarium and it also suffered severe losses.

"The younger Linné complained of the terrible damage done by mice, moulds and insects * * * [he] must have withdrawn the damaged sheets." (Jackson l. c. 21.) And we further learn from the son that Linnaeus himself destroyed many of his specimens. "My late father weeded out his herbarium, while he was able to work, and seems to have burned all the duplicates; why, no one knows." (Fries, Linné 2: 416, note.)

In short, in dealing with specimens now in the Linnaean herbarium, we are very frequently indeed dealing with specimens which were not specifically referred to by him in his works and which we cannot feel sure were in his possession at any particular date. All we can feel sure from his naming is that he referred a particular specimen to some particular species of his own. And unfortunately his knowledge of his own species was very frequently vague and unreliable. He was engaged in the herculean task of putting into usable form the works of his predecessors, and neither he nor anyone else under the circumstances could have been expected to have anything but the most general knowledge of the great mass of material with which he dealt.

Not infrequently, in dealing with one of the American species of Linnaeus we are confronted with the choice of applying a name given by him either (1) to a specimen of Clayton's collecting

which is definitely cited by Linnaeus and which we know he studied but which was not preserved in his own herbarium; or (2) to a specimen in the herbarium of Linnaeus but not cited by him, frequently without data and the history of which is entirely unknown, but one which bears his naming.

To test this matter let us suppose that an American author a number of years ago in a descriptive list of some collection not in his own herbarium, referred some particular specimen to some previously described species; then after a space of years let us suppose that in another work he gave a specific reference to his earlier publication and in this second work assigned a binomial name to such plant; let us further suppose that in his own herbarium at the time of the publication of his second work he had a sheet containing a specimen without data of any kind on which he wrote the binomial name given in his second work, but that he did not refer to such specimen in such second work; let us further suppose that this specimen represented a species other than the plant described in the first work. On such a state of facts I believe that all will agree that the binomial name in the author's second work should be applied to the plant described in the author's first work and should not be applied to the plant represented in his herbarium.

The above supposititious case represents a condition which is of frequent occurrence in dealing with Linnaean species of American plants, where he cites material collected by Clayton, and writers sadly led astray by the glamour of the Linnaean herbarium have reached results in identifying his species which would never have been thought of in other connections.

In such cases it seems to me that we should apply the Linnaean names to the specimens collected by Clayton; that we should follow certainly rather than uncertainly, definitely cited specimens rather than specimens merely named in an author's herbarium.

Let us apply the above to the following concrete cases:

(1) *PRUNUS VIRGINIANA* L. Sp. Pl. 473. 1753.

The original publication by Linnaeus reads as follows:

"2. *PRUNUS floribus racemosis, foliis deciduis basi antice glandulosis.*

Cerasus sylvestris, fructu nigricante in racemis longis pendulis phytolaccae instar congestis. Gron. virg. 54. Roy. lugdb. 537.

Cerasi similis arbuscula mariana, padi folio, flore albo parvo racemoso. Pluk. mant. 43. t. 339. Catesb. car. 1. p. 28. t. 28.

Habitat in Virginia."

The Clayton specimen cited from Gronovius is *Prunus serotina* Ehrh.

The only specimen in the Linnaean herbarium is *Prunus nana* DuRoi. A specimen was in the Linnaean herbarium in 1753. According to information furnished me by the late Dr. B. D. Jackson, there are no data of any kind whatsoever in connection with this specimen (a flowering one of which I have a photograph) to show who collected it or from where it came. It is entirely probable that it came from a cultivated plant in the Upsala Garden, because *Prunus virginiana* was listed as cultivated there in 1753 (Hojer, Dem. Pl. in Hort. Ups., 13), but this is only a supposition on my part.

The citation from Plukenet refers to *Itea virginica* L. and the citation from Catesby to *Prunus caroliniana* (Mill.) Ait. These references were cancelled by Linnaeus (Sp. Pl. Ed. 2) and can be disregarded. (See Torrey & Gray, Fl. N. Am. 1: 410.)

It will be noted that the Linnaean specific name was taken from the Clayton specimen, it being the only one cited from Virginia. It will be noted also that Linnaeus gave no description of his own, except such as is contained in his polynomial name. This applies to either *Prunus serotina* Ehrh. or to *Prunus nana* DuRoi; and as a matter of fact merely followed a system he adopted for naming various species of *Prunus*, his names running (1) "*Prunus floribus racemosis, foliis deciduis basi subtus biglandulosis*"; (2) "*Prunus floribus racemosis, foliis deciduis basi antice glandulosis*"; (3) "*Prunus floribus racemosis, foliis sempervirentibus eglandulosis*," etc.

Under these circumstances it seems to me that Miller, DuRoi, Wangenheim, Marshall, Aiton, Walter, Poiret, Pursh, Bigelow, Elliott and numerous more recent writers have been correct in applying the Linnaean name to the black cherry (*Prunus serotina* Ehrh.) and I cannot follow Prof. Fernald's contrary course based on partial information. (RHODORA 18: 140.)¹

(2) ASTER NOVAE-ANGLIAE L. Sp. Pl. 875. 1753.

The original publication by Linnaeus is as follows:

¹ It may be here noted that a strikingly similar problem is involved in *Quercus rubra* L. (Sp. Pl. 996). The specimens in the Linnaean Herbarium so named by Linnaeus are specimens of *Quercus coccinea* Wang. (Sargent in RHODORA 18: 45-6), and the Linnaean polynomial name applies to them as well as to other related species. However, they are not cited or referred to by him, altho they were apparently in his herbarium in 1753. Under these circumstances Sargent disregarded these specimens and applied the Linnaean name to the Clayton material which was directly cited by Linnaeus. It seems to me that in so doing he was quite correct. (See RHODORA 17: 39-40 and 18: 45-8.)

"15. ASTER foliis lanceolatis alternis integerrimis semiamplexicaulibus, floribus terminalibus. *Hort. cliff.* 408. *Gron. virg.* 100. *Roy. lugdb.* 166.

Aster novae angliae altissimus hirsutus, floribus amplis purpuro-violaceis. *Herm. par.* 98. *t.* 98.

Aster novae angliae altissimus hirsutus, floribus omnium maximis purpuro-violaceis. *Tournef. inst.* 482.

Habitat in Nova Anglia. 24

Caulis fuscus. Pedunculi imbricati foliolis. Corollae radius caeruleus."

The earlier citations given by Linnaeus refer to the plant commonly passing as *Aster novae-angliae* L.

The specimen in the Linnaean herbarium is of *Aster grandiflorus* L. (Gray, *Proc. Am. Acad.* 17: 165. 1882). A specimen (probably this) was in the Linnaean herbarium in 1753.

The eight words of description by Linnaeus apply to *Aster grandiflorus* L. and not to *Aster novae-angliae* L. as ordinarily understood.

Here we would undoubtedly have applied the Linnaean name to the plant which is represented in his herbarium and to which his few words of original description apply, were it not for the fact that he took his specific name from the plant of earlier authors, in fact one which he himself had dealt with in the Hortus Cliffortianus. But as far as I can see the case is much stronger for applying the name *Aster novae-angliae* to *Aster grandiflorus* than is the case for applying the name *Prunus virginiana* to the choke-cherry.

(3) ACALYPHA VIRGINICA L. Sp. Pl. 1003. 1753.

The original publication by Linnaeus reads as follows:

"1. ACALYPHA involucris femineis cordatis incis, foliis ovato-lanceolatis petiolo longioribus. *Hort. ups.* 290. *Fl. zeyl.* 342.

"Acalypha foliis ovato-lanceolatis, involucris femineis obtusis. *Hort. cliff.* 495. *Gron. virg.* 116.

"Mercurialis tricoccus hermaphroditica s. ad foliorum juncturas e foliolis cristatis julifera simul & fructum gerens. *Burm. zeyl.* 248. *t.* 99. *f.* 4. (Should be *Pluk. phyt.*)

"*Habitat in Zeylona, Virginia.* ☉"

Investigation has shown that the only specimen of this species in the Linnaean herbarium is without data of any kind. Of this specimen I have received photographs. It is highly probable that it was taken from the Upsala Garden, but this is at best only a guess. Mueller's definite statement that it came from the Upsala Garden referred to by Mr. Weatherby in his very thoughtful study of this species and its allies (RHODORA 29: 197) was not justified as far as

I have been able to find out. According to Weatherby (l. c. 196-7) this specimen represents one species, and the Clayton specimen from Virginia cited by Linnaeus represents another. Had we an authentic specimen from the Upsala Garden we would be confronted with the choice of applying the name of Linnaeus either to such specimen or to the specimen collected by Clayton from which he took his specific name. In that case it seems to me that the latter course would have been the correct one. But when in addition we find that there is no authentic material preserved from the Upsala Garden, it seems to me very plain that we must apply the Linnaean name to the Clayton specimen.

(4) *SCIRPUS CAPITATUS* L. Sp. Pl. 48. 1753.

The original publication by Linnaeus reads as follows:

"5. *SCIRPUS* culmo tereti nudo setiformi, spica subglobosa.

Scirpus culmo setaceo nudo, spica subglobosa. Gron. virg. 12.

Habitat in Virginia."

The Clayton specimen cited from Gronovius is *Eleocharis tenuis* (Willd.) Schultes.

The only specimen in the Linnaean herbarium is one of *Eleocharis obtusa* (Willd.) Schultes. This was one of those specimens "obtained after 1767, or * * by some accident not recorded by Linné" (Jackson).

Dr. Blake (*RHODORA* 20: 24) ignored the specimen in the Linnaean herbarium and applied the Linnaean name to the Clayton material. In this it seems to me that he was entirely correct.

MAPLEWOOD, NEW JERSEY.

THE GENUS TRISETUM IN AMERICA

FATHER LOUIS-MARIE O. C.

(Continued from page 222) 3

TRISETUM OREOPHILUM Louis-Marie, var. **Johnstonii**, var. n. A typo differt panícula cylindrica (3 cm. long., 1 cm. lat.), exserta; spiculis 3-floris; arista maxime variabili: a mucrone vix 1 mm. long. ad normalem 4 mm. long. divaricatam aristam; pallida canescentia laminarum vaginarumque.

Differing from the type by panicle (3 cm. long, 1 cm. wide) cylindrical, exserted; spikelets 3-flowered; awn very variable: from a short (less than 1 mm. long.) beak to a normal (4 mm. long) divaricate one; by the canescent pilosity of the blades and sheaths.

- Distribution. North ARGENTINA: Andes of N. W. San Juan, Arroyo Tambillos, 4300 m. alt. Jan. 10, 1926, *Ivan M. Johnston*. TYPE at the Gray Herbarium; isotype at the U. S. Nat. Herbarium of Washington.
22. *TRisetum LASIOLEPIS* E. Desv., C. Gay, Flor. Chil. **6**: 346. 1853.
23. *TRisetum PRESLEI* (Kunth) E. Desv. C. Gay, Flor. Chil. **6**: 347. 1853.
- TRisetum PRESLEI* (Kunth) E. Desv., var. **lasiantha** (Phil.), comb. n. *Deschampsia lasiantha* Phil., *Linnaea* **33**: 290. 1864-65.
- TRisetum PRESLEI* (Kunth) E. Desv., var. **Buchtienii** (Hack.) comb. n. *Trisetum Buchtienii* Hack. Oesterr. botan. Zeitschr. **54**: 290. 1904.
24. *TRisetum SPICATUM* (L.) Richt. Pl. Eur. **1**: 59. 1890. *Aira spicata* L. Sp. Plant. n. 7, 64, non n. 1. 63, 1753. *Aira sub-spicata* L. Syst. Nat. **2**: 873. 1759. *Trisetum subspicatum* Beauv., Nouv. Agrost. **88**. 1812. *Avena subspicata* Clairville, Man. Suisse, **17**. 1811. *Avena airoides* Koeler, Descr. gram. 205. 1902. *Trisetum airoides* Roemer & Schultes, Syst. Veg. **2**: 666. 1817. *Melica triflora* Bigel, New Engl. Journ. Med. et Surg. **5**: 334. 1816. White Mts. (fide Piper). *T. toluicense* Kunth, Rev. Gram. **1**: 101 & 297, t. 60. 1830 (1829). *T. andinum* Phil. *Linnaea* **29**: 93. 1857-58. non Bentham 1846. *T. labradoricum* Steud. Syn. Pl., Gram. 228. 1854. *T. groenlandicum* Steud. Syn. Pl., Gram. 228. 1854. *T. subspicatum* var. *breviglume* Hack. Wiss. Ergbn. Südpolar-Exp. **4**, pt. 4: 6. 1906. *T. subspicatum* var. *glabrifolium* Hackel, Rep. Princeton Univ. Exped. Patagonia, **8**, Suppl. 49. 1915.

One must go back to Scheuchzer,¹ quoted by Linnaeus in his original diagnosis of *Aira spicata*,² to reconstitute the type of *T. spicatum*, where the latter species is fully described and illustrated.

The plant (17-30 cm. high) is described as having glabrous, striate leaves; ligule (about 1 mm. long) obtuse; culm densely tomentose; panicle spike-like (about 1.5 cm. long, 0.7-1 cm. wide), purplish, shining; spikelets (about 4 mm. long) 2-flowered; glumes unequal (I shorter and narrower, II about 2 mm. wide and 4 mm. long), glabrous; lemma (4 mm. long) awned dorsally on the upper $\frac{1}{4}$; awn (about 4 mm. long) reflexed; palea equal or subequal to the lemma; rachilla densely villous; anthers 0.6-1 mm. long.

In the present state of our knowledge, the following variations departing from Scheuchzer's type seem to deserve mention:

¹ Scheuchzer, Johan., *Agrostographia*, 221. 1719.

² *Species Plantarum*, n. 7, 64, non n. 1, 63. 1753.

1. *T. SPICATUM* var. **hirsutum** (Phil.), comb. n. *T. hirsutum* Phil. Anal. Univers. Chil. 565. July 1873.
2. *T. SPICATUM* var. **PHLEOIDES** (Kunth) Macloskie, Rep. Princeton Univers. Exped. Patagonia 8: 206. 1904.
3. *T. SPICATUM* var. **fuegianum** (Hack.), comb. n. *T. subspicatum* var. *fuegianum* Hack. Wiss. Ergebn. Schwed. Südpolar-Exped. 4, pt. 4: 1906.
4. *T. SPICATUM* var. **dianthemum**, nom. n. *T. biflorum* Phil. Anal. Univ. Chil. 568. Jul. 1873; non Hochst. Flora 38: 275. 1852.
5. *T. SPICATUM* var. **andinum** (Benth.), comb. n. *T. andinum* Benth. Pl. Hartw. 261. 1847. *T. subspicatum* var. *compactum* Lange, Consp. Fl. Groenland, 3: 164. 1880.
6. *T. SPICATUM* var. **laxius** (Lange), comb. n. *T. subspicatum* var. *laxius* Lange, Consp. Fl. Groenl. 3: 164. 1880.
7. *T. SPICATUM* var. **nivosum** (Fourn.), comb. n. *T. nivosum* Fourn., Mex. Pl. Enum., Gram. 107. 1886.
8. *T. SPICATUM* var. **MAJUS** Farwell. Rep. Mich. Acad. Sci. 21: 352. 1920.
9. *T. SPICATUM* var. **MOLLE** (Michx.) Piper, Contr. U. S. Nat. Herb. 11: 125. 1906.
10. *T. SPICATUM* var. **Brittonii** (Nash), comb. n. *T. Brittonii* Nash, Bull. N. Y. Gard. 1: 437. 1900.
11. *T. SPICATUM* var. **PILOSIGLUME** Fern. RHODORA 18: 195. 1916.
12. *T. SPICATUM* var. **MAIDENII** (Gand.) Fern. RHOD. 18: 196. 1916.
13. *T. SPICATUM* var. **alaskanum** (Nash) Malte ined. *T. alaskanum* Nash Bull. N. Y. Botan. Garden 2: 155. 1901.
14. *T. SPICATUM* var. **villosissimum** (Lange), comb. n. *T. subspicatum* var. *villosissimum* Lange, Consp. Fl. Groenl. 3: 164. 1880.

H. Glumes very unequal. . . I.

I. Culm pubescent or puberulent under the panicle;
palea 2-toothed.

25. *TRISETUM FOURNIERANUM* Hitchc. Contr. U. S. Nat. Herb. 17: 326. 1913.
26. *TRISETUM ROSEI* Scribn. & Merrill, Contr. U. S. Nat. Herb. 8: 289. 1905.
TRISETUM ROSEI Scribn. forma **tenerum** (Scribn. & Merrill), comb. n. *T. Rosei* var. *tenerum* Scribn. & Merrill, Contr. U. S. Nat. Herb. 8: 289. 1905.
27. *TRISETUM CONGDONI* Scribn. & Merrill, Bull. Torr. Bot. Club 29: 470. 1902.

Trisetum Congdoni is easily distinguished from *T. spicatum* by its spikelets longer, silvery shining, with glumes (6 and 8 mm. long in the type) narrow and long-attenuated at the tip, with lemma (7-8 mm.) long-awned; by its leaves more erect and glabrous.

This species is neither to be confused with *T. sesquiflorum* Trin., as

pointed out elsewhere; the latter being of boreal distribution, with more narrow-lanceolate glumes, with awn attached dorsally near the base of the lemma.¹

28. *TRisetum* *barbinode* Trin. *Linnaea* **10**: 300. 1836.

TRisetum *barbinode* var. *hirtiflorum* (Hack.), comb. n. *T. hirtiflorum* Hack. Fedde Rep. Spec. Nov. **10**: 169. 1911.

I. *Culm glabrous under the panicle; palea 2-fid. . . J.*

J. *Anthers 0.6-1 mm. long.*

29. *TRisetum* *caudulatum* Trin. *Linnaea* **10**: 300. 1836. *T. chromostachyum* E. Desv., C. Gay Fl. Chil. **6**: 350. 1853.

30. *TRisetum* *variabile* E. Desv., C. Gay. Fl. Chil. **6**: 351. 1853.

1. *T. VARIABILE* var. a) *FLAVESCENS* E. Desv. *T. ochrostachyum* Phil. *Linnaea* **33**: 290. 1864-65.

2. *T. VARIABILE* var. b) *VIRESCENS* E. Desv.

3. *T. VARIABILE* var. *intonsum* nom. n. *T. variabile* var. *virescens* (Nees) Mackloskie, not E. Desv.

4. *T. VARIABILE* var. *chiloense* (Phil.), comb. n. *T. chiloense* Phil. *Linnaea* **29**: 93. 1857-58.

5. *T. VARIABILE* var. *Vidali* (Phil.), comb. n. *T. Vidali* Phil. Anal. Univ. Chil. **94**: 26. 1896.

31. *TRisetum* *erectum* Phil. Anal. Univ. Chil. **94**: 26. 1896.

J. *Anthers 1-2 mm. long.*

32. *TRisetum* *monticola* Phil. *Linnaea* **33**: 291. 1864-65.

33. *TRisetum* *antarcticum* Trin. Mém. Acad. St. Pétersb. Sér. 6. **1**: 61. 1830.

A. *Panicles axillary; culm branching at each node.*

34. *TRisetum* *paradoxum* Phil., Anal. Univ. Chil. **94**: 28. 1896.

Sub-section 2. SPHENOPHOIDEA Louis-Marie.

35. *TRisetum* *pennsylvanicum* Beauv. in R. & S. Syst. **2**: 658. 1817.

36. *TRisetum* *interruptum* Buckl. Proc. Acad. Sci. Phila. **100**. 1862.

TRisetum *interruptum* Buckl., var. *californicum* (Vasey), comb. n. *T. californicum* Vasey, U. S. Dept. Agr. Div. Bot. Bull. **12**: Pl. 46. 1892.

37. *TRisetum* *hallii* Scribn. Bull. Torr. Bot. Club **2**: 6. 1884.

Sub-section 3. GRAPHEPHORUM (Desv.) Louis-Marie.

38. *TRisetum* *melicoides* (Michx.) Vasey, Bot. Gaz. **9**: 169. 1884.

39. *TRisetum* *wolfii* Vasey, Monthly Report U. S. Dept. Agr. Mar. **156**. 1874.

¹ Abrams, *Flora of the Pacific States* **1**: 168 (1923) by error gives *T. Congdoni* as a synonym to *T. sesquiflorum*. Prof. A. S. Hitchcock wrote us to this effect.

TRISETUM WOLFII Vasey, var. **Brandegei** (Scribn.), comb. n. *T. Brandegei* Scribn. Bull. Torr. Club. **10**: 64. 1883. *Grappophorum Brandegei* (Scribn.) Rydb. Rocky Mount. Fl. 61. 1917.

TRISETUM WOLFII Vasey, var. **BRANDEGEI** (Scribn.) Louis-Marie, forma **muticum** (Boland.), comb. n. *T. subspicatum* var. *muticum* Boland. in S. Watson, Bot. Calif. **2**: 296. 1880. *Trisetum muticum* Scribn. U. S. Dept. Agr. Div. Agrost. Bull. **11**: 50 f. 10. 1898. *T. Wolfii* Vasey, var. *muticum* Scribn. RHODORA **8**: 88. 1906.

40. TRISETUM ALTIJUGUM (Fourn.) Scribn. RHODORA **8**: 89. 1906.

41. TRISETUM PRINGLEI (Scribn.) Hitchc. Proc. Biol. Soc. Wash. **40**: 82. 1927.

Sub-section 4. KOELERIFORMIA Louis-Marie.

Discussion. In 1805, Persoon published, side by side, in his Synopsis, Trisetum and Koeleria;¹ he thus characterized the latter one: "173. KOELERIA. *Cal.* multiflorus, 2-valvis, compresso-carinatus. *Cor.* 2-valvis, brevi-aristata: glumis nervosis. (Spica composita ex spiculis compressis, saepius pubescentibus subsessilibus.)" "173. KOELERIA *Cal.* many flowered, 2 valved, compressed-keeled. *Cor.* 2-valved, short-awned: glumes nerved (spike composed of compressed spikelets, more often pubescent, sub-sessile)."

The distinction, really weak, between the two genera, is reduced to " . . . *Cor.* short-awned (*Cor.* brevi-aristata)." The capricious historical development, fickle and unsteady, of the conception of the two genera in question, weakening still the line of demarcation, makes today the classification of the extreme species neighboring *Koeleria* and *Trisetum* very hard, often very subjective.

"Le genre *Koeleria*," says E. Cosson, "ne se distingue du genre *Trisetum* que par le caryopse muni d'un sillon et velu, et non pas dépourvu de sillon et glabre. Malgré le peu de valeur de ces caractères différentiels, nous avons été amenés à séparer génériquement les *Trisetum* des *Avena*, car, ainsi que M. Émile Desvaux (*Gramineae Chilenses*) l'a très bien reconnu, les *Trisetum* sont encore plus distincts des *Avena* qu'ils ne le sont des *Koeleria*." ²

"It is difficult," says Beal,³ "to assign to *Koeleria* any positive character. The glumes are more scarious and have fainter nerves than in the others of the subtribe."

The difficulty met in separating the two genera in question is

¹ Persoon, C., Synopsis Plantarum **1**: 97. 1805.

² Cosson, E., Bull. Soc. Bot. France **1**: 12. may. 1854.

³ Beal, W. J., Grasses of N. Amer. **2**: 495. 1896.

not such as should surprise us too much. The *Aveneae* and *Festuceae* groups, whose extreme types are so clean cut, just blend into one another by numerous intermediary species; their principal characteristics, taken from the length of the glumes, the insertion of the awn and its form, are far from always being sufficiently distinctive. Such is, for example, the genus *Koeleria* which is hardly separated from the genus *Trisetum* and by that very fact belongs to the tribe of the *Aveneae*, and which may be equally referred to the tribe *Festuceae*, where most authors class it.¹

42. TRisetum LAXIFLORUM Phil. Anal. Univ. Chil., 567. Jul. 1873.
43. TRisetum ARAEANTHUM Phil. Anal. Univ. Chil., 94: 26. 1896.
44. TRisetum BRACHYATHERUM Phil. Anal. Univ. Chil., 567. Jul. 1873.
45. TRisetum DEPAUPERATUM Phil. Linnaea 33: 291. 1864-65.
46. TRisetum MICRATHERUM E. Desv., C. Gay, Flora Chil. 6: 352. 1853.
47. TRisetum NEMOROSUM Phil. Anal. Univ. Chil. 94: 26. 1896.
48. TRisetum LAXUM Phil. Anal. Univ. Chil. 568. 1873.

Sub-section 5. DESCHAMPSIOIDEA Louis-Marie.

49. TRisetum PALMERI Hitchc. Contrib. U. S. Nat. Herb. 17: 325. 1913.
50. TRisetum **brasiliense**, sp. n. Rigidum, dense caespitosum (35-50 cm. altum), laeve et glabrum; culmis strictis, filiformibus; vaginis striatis, brevibus, marcescentibus; ligula (minor 1 mm.) bilateraliter producta in dua auricula, laminam lateraliter excedentia, centrali parte depressa; laminis foliorum angustioribus quam vaginae, involuto-filiformibus: his fasciculorum sterilium facile 25 cm. longis, his caulinis brevioribus, saepe ad oram vaginae fractis; panícula (5-8 cm. long., 1 cm. lat.) pauciflora, straminea-flava; spiculis bifloris: floribus plus minusve caducis; glumis aequilongis, 3-nerviis saltem inferius, navicularibus, glabris, moderate acuminatis, spiculam aequantibus: I (1.5 mm. lat.), II (2 mm. lat.) magis amplectante; lemmate (4.5 mm. long., 2 mm. lat.) apice bifido (1 mm.) segmentis latis, acutis, 5-nervio, nervis crassis, supra (0.5 mm.) basim aristato; arista (5-6 mm. long.) dimidia parte inferiore tortili vel rarissime recta, superius geniculata vel diverso recurvata; interfloris articulis rachillae et ejus prolongatione (0.5-0.7 mm. long.) perbrevibus; callo albo-piloso; palea lemma subaequante, carinis (0.6 mm. distantibus) dense pilosis; antheris (2 mm. long.) linearibus.

Culm rigid, densely cespitose (35-50 cm. high), smooth and glabrous, filiform; sheaths striate, short, marcescent; ligule (less than 1 mm. long) reduced in the centre, auriculate

¹ Cosson, E., loc. cit. 1854.

laterally; blades narrower than the sheaths, involute-filiform: the sterile shoots of blades easily 25 cm. long, the cauline ones shorter, breaking up easily at the base; panicle (5–8 cm. long, 1 cm. wide) few-flowered, stramineous-yellow; spikelets 2-flowered; glumes equally long, 3-nerved, navicular, glabrous, acuminate, equal to the spikelet: I (1.5 mm. wide), II (2 mm. wide) more inclosing; lemma 4.5 mm. long., 2 mm. wide) 2-fid (1 mm. long) at the apex, 5-nerved, awned near the base; awn (5–6 mm. long) strongly twisted at the inferior half, geniculate or diversely bent at the superior half; rachilla and callus covered with white hair; palea subequal to the lemma; keels (0.6 mm. distant) densely pilose; anthers (2 mm. long) linear.

Distribution. BRAZIL: dense tufts in peaty soil among rocks above timberland, Itatiaia, Rio de Janeiro, Alt. 2200–2400 m., Jan. 17, 1925, *Agnes Chase*. 8304. TYPE in the U. S. Nat. Herb. Co-types at the Gray Herbarium and the Oka Agricultural Institute Herbarium. The Herbariums of Vienna and Brazil will receive each a share of the type.

Remark. We are indebted to Mrs. Agnes Chase's kindness for the above new species, which clearly belongs to the subsection *Deschampsioidea*. In fact, it was first classified under *Aira*.

51. *TRisetum JUNGENSEI* Hack. Verhandl. Zool. Bot. Gesellsch. Wien **65**: 75. 1915.

Section 2. *AULACOA* Louis-Marie.

52. *TRisetum Trinii* (Desv.), comb. n. *Bromus Trinii* Desv., C. Gay, *Flora Chilena* **6**: 441. 1853. *Trisetum barbatum* Steud. Syn. Pl., Gram. 229. 1854. *Bromus barbatoides* Beal, *Grasses N. Amer.* **2**: 615. 1896; Vasey. *Illust. N. Amer. Grasses II.* **2**: Plate 60. 1893.
1. Var. **pallidiflorum** (Desv.), comb. n. *Bromus Trinii*, var. α *pallidiflora* Desv. in C. Gay, *Fl. Chil.* **6**: 441. 1853.
 2. Var. **micrantherum** (Desv.), comb. n. *Bromus Trinii*, var. β *micranthera* Desv. l. c. 1853.
 3. Var. **manicatum** (Desv.), comb. n. *Bromus Trinii*, var. γ *manicata* Desv. l. c. 1853.
 4. Var. **effusum** (Desv.), comb. n. *Bromus Trinii*, var. δ *effusa* Desv. l. c. 442. 1853.
 5. Var. **strictum** (Desv.), comb. n. *Bromus Trinii*, var. ϵ *stricta* Desv. l. c. 1853.
 6. Var. **majus** (Vasey), comb. n. *T. barbatum* var. *major* Vasey, *Illust. N. Amer. Grasses II.* **2**: 1893. *Bromus barbatoides* var. *sulcatus* Beal, *Grasses N. Amer.* **2**: 615. 1896.
 7. Var. **litorale** (Phil.), comb. n. *T. litorale* Phil. *Linnaea* **29**: 92. 1857–58.
53. *TRisetum FLORIBUNDUM* Pilger, Bot. Jahrb. Engler, **37**: 505. 1906.

TRISETUM FLORIBUNDUM var. **Weberbaueri**, comb. n. *T. Weberbaueri* Pilger, Bot. Jahrb. Engler. **37**: 506. 1906. (*Weberbauer* 3078).

Sub-genus II. **ISOLYTRUM** Louis-Marie.

Deyceuxia Clar. (a Fourn. emend. Bull. Soc. Bot. Fr. **24**: 181. 1877) ut genus, in part.

54. TRISETUM **humile**, nom. n. *Deyceuxia gracilis* Fourn. Mex. Pl. Enum. Gram. 106. 1886; non H. A. Weddell, Bull. Soc. Bot. France. **22**: 179. mai 1875.
55. TRISETUM DEYEUXIOIDES (H. B. K.) Kunth. Rev. Gram. **1**: 102. 1829.
56. TRISETUM VIRIDE (H. B. K.) Kunth. Rev. Gram. **1**: 101. 1829.
57. TRISETUM LONGIGLUME Hack. Fedde. Rep. Nov. Spec. **7**: 319. 1909.
58. TRISETUM **andicola**, sp. n. Planta (20–30 cm. alta) robusta, caespitosa; ligula (1.5–2 mm. long.) lacerata; foliorum laminis (8–10 cm. long., 1.5–2 cm. lat.) glabris, dorsaliter subscabris; panicula (5 cm. long, 1.7 cm. lat.) spiciformi-densa, ovalilanceolata, ascendentibus ramis; spiculis bifloris; glumis 3-nerviis saltem inferius, aequilongis (8–9 mm. long.), subaequilatis: I (1.5 mm. lat.), II (1.6 mm. lat.), glabris, spiculam aequantibus; lemma (5–6 mm. circ. long., 1.8–2.2 mm. lat.) apice (1 mm.) bilobo, bisetulato, lobis duobus lateraliter scariosolaciniatis; ad 2.5 mm. e basi aristato; arista tenui (8 mm. circ. long.) recta vel saepius geniculata, infra genu distans 2.5–3 mm. ab insertionis loco, vix torta; rachilla et ejus pedicelliformi prolongatione longissime (3–4 mm.) albo-villosis, flores aequantibus; palea (5–5.5 mm. long.) carinis 0.3 mm. circ. distantibus; antheris (0.8–1 mm. long.) crassis.

Culm strong (20–30 cm. high) cespitose; ligule lacerated (1.5–2 mm. long.); blades glabrous (8–10 cm. long, 1.5–2 cm. wide) lightly scabrous on the ventral side; panicle (5 cm. long, 1.7 cm. wide) spiciform-dense, ovoid-elongate, with ascending branches; spikelets 2-flowered; glumes nearly isomorphous, 3-nerved at least at the base: I (1.5 mm. wide), II (1.6 mm. wide) glabrous, equal in size to the spikelets; lemma (5–6 mm. long, 1.8–2.2 mm. wide) very 2-lobed: lobes 2-setulate, scarious-lacinate on the sides; awn (8 mm. long) inserted at 2.5 mm. from the base, straight or more often geniculate at 2.5–3 mm. from the insertion point, lightly twisted under the bend; rachilla and the pedicelliform extension very (3–4 mm.) villous: white hairs equalling the glumes; palea (5–5.5 mm. long) equalling the lemma in size, 2-keeled: keels about 0.3 mm. distant; anthers (0.8–1 mm. long) broad.

Distribution. CHILI: Laguna Negro, 2700–4000 m. alt. Martio 1873. *Fr. Vidal Gorus* 265.

Discussion. Philippi never knew just where to place the plant. *Arundo?* *Danthonia?* The names he gave it—we are relieving the synonymy of such—prove the case. On examination its place is not dubious. On an herbarium sheet of Hackel, I find a manuscript note where the present combination is already proposed.

59. *TRISETUM EVOLUTUM* (Fourn.) Hitchc. Contrib. U. S. Nat. Herb. **17**: 325. 1913.

60. *TRISETUM MACBRIDEI* Hitchc. Contrib. U. S. Nat. Herb. **24**: 8. 359. 1927.

Before concluding, the writer takes great pleasure in thanking Prof. M. L. Fernald, under whose highly qualified guidance the work was accomplished, and the competent personnel of the Gray Herbarium. He acknowledges thanks also to the curators of the U. S. Nat. Herb. of Washington, of the N. Y. Bot. Garden, of the University of California Herbarium and of the National Herbarium of Canada. For the loan of material belonging to private collections, he thanks especially the RR. Brothers Marie-Victorin and Rolland-Germain and Prof. Lorenzo Parodi. He finally acknowledges thankfully the interest shown by Prof. A. S. Hitchcock and Mrs. Agnes Chase during his stay in Washington.

OKA AGRICULTURAL COLLEGE,

La Trappe, Quebec.

Vol. 30, no. 359, including pages 209 to 228, was issued 17 December, 1928.

ERRATA

- Page 3, line 26, for RUBRA read RUBRUM
 " 4, " 8, for GROELANDICUM read GROENLANDICUM
 " 4, last line, for GYMNSTOMUN read GYMNSTOMUM
 " 5, line 32, for FASICULARE read FASCICULARE
 " 8, " 13, for SUBGLOSUM read SUBGLOBOSUM
 " 10, " 18, for PLUMULOSUM read PLUMULOSUS
 " 16, " 10, for *Miss* read *Mrs.*
 " 48, " 13, for *gaspensis* read *gaspense*
 " 85, " 14, for 78 read 77
 " 99, " 27, for *philogenetic* read *phylogenetic*
 " 108, lines 3-4, for road, which crosses the Neponset River meadows at the station known as Dedham Road, read road (Neponset Street)
 " 121, line 19, for *gemmiscapa* read *geminiscapa*
 " 133, " 10, for Lunelll read Lunell
 " 136, " 8, for *guadaloupensis* read *guadalupensis*
 " 171, " 33, for *specmen*, read *specimen* s
 " 172, " 19, for *lateriflorus* read *lateriflora*
 " 176, last line, for " " "
 " 186, line 34, for *northeastern* read *northwestern*
 " 198, " 35, for *trifrida* read *trifida*

INDEX TO VOLUME 30.

New scientific names are printed in full-face type.

- Abietinella abietina*, 9
Acalypha virginica, 236
Acrothrix, 195
Actaea spicata, 71
 Additions to the Flora of Cape Cod,
 135; to the Flora of Worcester
 County, Mass., 20
Adiantum, 13; *pedatum*, 13
Agoseris, 77
Ahnfeldtia plicata, 196
Aira, 243; *spicata*, 238; *subspicata*,
 238
Alisma, 55
Amberboi, 59
Amblyodon dealbatus, 8
Amblystegiaceae, 9
Amblystegiella Sprucei, 10
Amblystegium serpens, 10
America, Genus *Oxytropis* in north-
 eastern, 137; Genus *Trisetum* in,
 209, 237; *Primula* § *Farinosae* in,
 59, 85
 American and eastern Asiatic Beck-
 mannia, The, 24; Forms of *Senecio*,
 Some eastern, 224; Occur-
 rence of *Athyrium alpestre*, The
 eastern, 44; Representatives of
 Asplenium Ruta-muraria, 37;
 Representative of *Calamagrostis*
 Epigejos, 202; Species and Vari-
 eties of *Rosa*, Ten New, 109;
 Variety of *Polystichum Braunii*,
 The eastern, 28
Ammophila, 202; *arenaria*, 204;
 breviligulata, 202-205
Anaulacoa, sect. of *Trisetum*, 211,
 212
Andracea alpestris, 2
Andraceaceae, 2
Andropogoneae, 215
Anemonanthea, sect. of *Anemone*,
 180, 181, 182; North American
 Species of, 180
Anemonanthea, 186; *quinquefolia*,
 186
Anemone, 180; § *Anemonanthea*,
 180-182, The North American
 Species of, 180; *cuneifolia*, 183;
 cyanea, 186; *deltoides*, 182, 188;
 Grayi, 180, 182, 186; *lancifolia*,
 181-185; *Lyallii*, 182, 187; *min-*
 ima, 181, 182, 185; *nemorosa*, 180,
 182, 183, 186, 188, β *quinque-*
 folia, 186, f. *quinquefolia*, 186,
 var. *quinquefolia*, 180, 182, subsp.
 americana, 186, var. *Lyallii*, 187,
 var. *oregana*, 186, 187, var. γ
 quinquefolia, 186; *oregana*, 182,
 186-188; *pedata*, 186; *Piperi*, 180,
 182, 187, 188; *quinquefolia*, 180,
 182, 185, 186, 188, var. *Grayi*,
 187, var. *Lyallii*, 187, var. *ore-*
 gana, 186; *trifolia*, 181, 183-185
Anisothecium rubrum, 3
Anomobryum concinnatum, 6
Antennaria, 77, 137
Aphanocapsa pulchra, 195
Aragallus arcticus, 144; *Belli*, 150,
 151; *deflexus*, 140; *foliolosus*, 140;
 hudsonicus, 142; *inflatus*, 154;
 johannensis, 146; *podocarpus*, 154
Arnica, 137
Arundo, 202, 245
 Asiatic Beckmannia, The American
 and eastern, 24
Asparagopsis, 198; *hamifera*, 197
Aspidium, 16; *aculeatum*, 30, var.
 Braunii, 30; *alpestre*, 48; *Boottii*,
 17; *cristatum*, 17, var. *Clintoni-*
 anum, 17; *Goldianum*, 17; *mar-*
 ginale, 17, f. *Davenportii*, 17, var.
 elegans, 17; *novboracense*, 16;
 simulatum, 16; *spinulosum*, 17,
 var. *concordianum*, 17, var. *dila-*
 tatum, f. *anadenium*, 17, var.
 intermedium, 17; *Thelypteris*, 16,
 f. *Pufferae*, 16
Asplenium, 15; **cryptolepis**, 41-43,
 var. **ohionis**, 43; *ebeneum*, f.
 Hortoniae, 14, var. *Hortoniae*, 14,
 var. *incisum*, 14; *ebenoides*, 14;
 Filix-femina, 15; *platyneuron*, 14,
 f. **Hortoniae**, 14, f. *serratum*, 14,
 var. *incisum*, 14; *Ruta-muraria*,
 14, 37-42, The American Repre-
 sentatives of, 37, var. *lanco-*
 latum, 43; *Trichomanes*, 14
Aster, 77, Notes on, 226; *bifrons*,
 178, 179; *columbianus*, 227; *cordi-*
 folius, 175; *coridifolius*, 165, 169;
 cyaneus, α *glauca*, 228; *diffusus*,
 172, 176, 178, var. *bifrons*, 175,
 178, var. *hirsuticaulis*, 172, 176,
 var. *horizontalis*, 175, var. *thy-*
 soideus, 175, 177, 178, var. *vari-*
 folius, 176; *divergens*, 173; *du-*

mosus, 162, 163, 165-167, 169, 171, 175, 176, \times *vimineus*, 172, var. *coridifolius*, 163, 165, 169, var. *Dodgei*, 164, 167, var. *gracilentus*, 166, 169; var. ***gracilipes***, 163, 166, var. ***pergracilis***, 163, 166, 169, var. *strictior*, 164, 165, 167, 169, var. *subulaefolius*, 164, 167-169, var. *subracemosus*, 172; *Engelmianii*, 228, var. *glaucescens*, 228; *ericoides*, 78, 171, 172, 175, \times *vimineus*, var. *subdumosus*, 172, var. *platyphyllus*, 106, in Maryland, 106, var. *villosus*, 79; *exiguus*, 227; *foliolosus*, 168, 172; *fragilis*, 165; *furcatus*, 77; ***glaucescens***, 228; *glaucophyllus*, 228; ***Gormanii***, 228; *grandiflorus*, 236; *hirsuticaulis*, 172, 176; *horizontalis*, 173, 175; *laevis*, 228; *lateriflorus*, 161, 162, 164, 171, 172, 175-179, and some of its Relatives, 161, var. ***angustifolius***, 164, 174-176, var. *bifrons*, 178, var. *glomerellus*, 172, 175, 177, var. *grandis*, 178, var. *hirsuticaulis*, 172, 176, var. *pendulus*, 164, 172, 173, 175, 176, var. ***tenuipes***, 164, 174, 175, var. *thyrsoides*, 177, 178; *ledophyllus*, 228; *miser*, 172, var. *diffusus*, 172, var. *glomerellus*, 172, 175, 177, var. *hirsuticaulis*, 172, 176; *missouriensis*, 165, 171, 175, 177-179, var. ***thyrsoides***, 165, 175, 177, 178, \times *vimineus*, var. *subdumosus*, 172; *multiflorus*, 78, 79, 175, 227, 228, A New Variety of, 77, var. ***caeruleus***, 78, var. ***columbianus***, 227, var. *exiguus*, 227, var. ***pansus***, 227; *memoralis*, 136; *novae-angliae*, 235, 236, var. *roseus*, 78; *novi-belgii*, 171; ***ontarionis***, 165, 175, 179; *paniculatus*, 161, 162, 171, 172, 175, 177-179; *paucicapitatus*, 228; *pendulus*, 173, 176; *pumiceus*, 175; *racemosus*, 172; *recurvatus*, 172; *salicifolius*, 78; *saxatilis*, 172, 175, 179; *secundiflorus*, 169; *Shortii*, 78; *sparsiflorus*, 166, 169; *spectabilis*, 226, var. ***cinerascens***, 226; *stricticaulis*, 227; *tataricus*, 79; *thyrsoides*, 175, 177, 178; *Tradescanti*, 170, 172, 178, 179, var. *fragilis*, 170; *undulatus*, 171, 175; ***vialis***, 228; *vimineus*, 162, 164, 169, 171, 172, 175, 176, 179, var. ***dubius***, 164,

170, 171, var. *foliolosus*, 171, 172, var. *saxatilis*, 172, var. ***subdumosus***, 164, 171, 172, 178

Asteraceae, 226

Asters, Some Illinois, and a New Variety of *A. multiflorus*, 77

Astragalus, 137; *aboriginum*, 122; *biflorus*, 154, 155; *deflexus*, 140; *leucanthus*, 142; *nigrescens*, 153; *pygmaeus*, 153; *retroflexus*, 140

Athyrium, 15; *acrostichoides*, 15; *alpestre*, 44, 47-49, The eastern American Occurrence of, 44, var. *americanum*, 44-49, var. ***gas-pense***, 48, 49, var. ***typicum***, 48; *americanum*, 44-47, 49; *angustifolium*, 15; *angustum*, 15, f. *confertum*, 15, f. *elegans*, 15, f. *laciniatum*, 15, var. *elatus*, 15, var. *rubellum*, 15; *asplenoides*, 15, f. *subtripinnatum*, 15; *Filix-femina*, 45, 48

Aulacoa, sect. of *Trisetum*, 211, 243

Aulacomniaceae, 8

Aulacomnium palustre, 8

Authors of *Sonchus arvensis*, var. *glabrescens*, 19

Avena, 241; *airoides*, 238; *subspicata*, 238

Aveneae, 242

Banisteria, 54

Barbula vinealis, 5

Bartram, E. B., Newfoundland Mosses collected by Mr. Bayard Long in 1924-26, 1

Bartramia ithyphylla, 9; *pomiformis*, 9

Bartramiaceae, 8

Beckmannia, 24-26, The American and eastern Asiatic, 24; *baikalensis*, 25-27; *erucaeformis*, 24-27, 51, var. *baikalensis*, 25-27, var. *uniflora*, 27, var. *uniflorus*, 26, 27; ***Syzigachne***, 27

Beech Fern, 16

Benke, H. C., Some Illinois Asters and a New Variety of *A. multiflorus*, 77; Two New Varieties of early Spring Plants—a *Ranunculus* from Missouri and a *Heterotheca* from Texas, 200

Betula, 125; *lutea*, 125, 129, \times *pumila*, 125, \times *pumila*, var. *glandulifera*, 125, var. *typica*, 129; *papyrifera*, 125, 126, 128; *pumila*, 125, 128, var. *glandulifera*, 125, 127, 128; *Purpusii*, 125, 126, 129, var. *luteoides*, 129;

- Sandbergi, 125-129, Evidence of the Hybrid Nature of, 125
- Bidens, 31; bidentoides, 31; cernua, 31; comosa, 31; connata, 31, 32, 34, Varieties of, in Wisconsin, 31, var. **ambiversa**, 31, 33, 35, var. *anomala*, 31, 34, var. *fallax*, 31, 32, var. *fultior*, 32, var. *petiolata*, 31-33, 35, var. *pinnata*, 31, 34, 35, var. **typica**, 31-33, 35; coronata, 34; Eatoni, 31; heterodoxa, 31; hyperborea, 31; laevis, 31; multiceps, 31; petiolata, 33; trichosperma, 34; tripartita, 31
- Bignonia, 58; capreolata, 58; radicans, 58
- Bladder Fern, 18
- Blairia, 54
- Blake, S. F., *Aster ericoides* var. *platyphyllus* in Maryland, 106; Notes on *Aster*, 226; *Potentilla intermedia* in the Boston District, 107
- Bonnemaisoniaceae, 197
- Boston District, *Potentilla intermedia* in the, 107
- Botrychium ramosum, 198; simplex, 198
- Brachylobus hispidus, 132, 133
- Brachytheciaceae, 10
- Brachythecium reflexum, 11, salebrosus, 11
- Bracken, 13
- Brake, 13; Cliff, 14; Rock, 14
- Breidleria arcuata, 11
- Bromus barbatooides, 243, var. *sulcatus*, 243; Trinii, 243, var. δ *effusa*, 243, var. γ *manicata*, 243, var. β *micranthera*, 243, var. α *pallidiflora*, 243, var. *stricta*, 243
- Brotherella delicatula, 11
- Bryaceae, 6
- Bryum affine, 7; argenteum, 8; bimum, 8; caespiticum, 8; calophyllum, 6; capillare, 7, var. *obconicum*, 8; globosum, 7; inclinatum, 6; Knowltoni, 6; **Longii**, 1, 2, 6; pallens, 7; pendulum, 6; ventricosum, 8
- Cakile edentula, 204; maritima, 204
- Calamagrostis, 202, 203; **arenicola**, 203-205; cinnoides, 203; Cummingii, 212; epigejos, 202, 203, An American Representative of, 202; Lechleri, 212; Pseudophragmites, 202; purpurascens, 219
- Calcitrapa, 59
- Calcitrapae, 59
- Calcitrapoides, 59
- Calliergon giganteum, 10; sarmenosum, 10; stramineum, 10; turgescens, 10
- Calluna vulgaris, 204, 205
- Camptosorus, 15, 42; rhizophyllum, 14, 15, f. *auriculatus*, 15
- Campylium stellatum, 10
- Campylophyllum Halleri, 10
- Cape Cod, Additions to the Flora of, 135
- Capriola Dactylon, 51
- Carex, 199; alba, 29; crinita, 199; eburnea, 29; gynandra, 199; incurva, 77; loliacea, 29; Macloviana, 77; Mitchelliana, 198-199, and other rare Plants near Cohasset, Mass., 198; trisperma, 29
- Catosciopiceae, 8
- Catoscopium nigrum, 8
- Centaurea, 59
- Ceratodon purpureus, 3
- Chain Fern, 14
- Chara, 195
- Characiopsis crassiapex, 194; **pileata**, 194
- Chenopodium Botrys, 158; carinatum, 158, and other unusual Weeds, 158
- Chondrus, 198; crispus, 196
- Christmas Fern, 16
- Chroococcus minutus, 195
- Churchill, J. R., Reports on the Flora of Massachusetts, I, 12
- Cinclidium stygium, 8
- Clandestine Form of *Leersia oryzoides*, The, 81
- Clayton, concerning the proper Identification of Linnaean Species based on Material collected by, 232
- Cliff Brake, 14
- Climaciaceae, 9
- Climacium dendroides, 9
- Coelosphaerium Naegelianum, 195
- Collins, J. F., Notes on certain Species of *Panicum* occurring in or near Rhode Island, 229
- Comandra, 21, 22; californica, 21; elegans, 21; livida, 21-24; pallida, 21; Richardsiana, 21; umbellata, 21
- Concerning the proper Identification of Linnaean Species, especially those Based on Material collected by Clayton, 232
- Contributions from the Gray Herbarium, LXXIX, 21, 37, 59, 85; LXXXII, 180

Convallaria Keiskei, 184; majalis, 183, 184; majuscula, 184
 Copeland, J. J., Characiopsis pileata, n. sp., 194
 Corallorhiza trifida, 198
 Corema alba, 204; Conradii, 204
 Corydalis sempervirens, 36; Exceptional Dimensions in, 36
 Cratoneurum filicinum, 9; glaucum, var. falcatum, 9
 Crocodilium, 59
 Crocodiloidea, 59
 Cryptogramma, 14; Stelleri, 14
 Cyani, 59
 Cyanus, 59
 Cynodontium polycarpum, 3
 Cypripedium Calceolus, 47; parviflorum, 47, var. planipetalum, 47
 Cystopteris, 18; bulbifera, 18; fragilis, 18

Dactylis, 50, 51
 Damasonium, 55
 Danthonia, 245
 Darbya, 22
 Deane, W., Reports on the Flora of Massachusetts, I, 12
 Dennstaedtia, 18; punctilobula, 18
 Deschampsia lasiantha, 238
 Deschampsioidea, sub-sect. of Trisetum, 211, 242, 243
 Desmatodon cernuus, 5; systylius, 1, 5
 Deyeuxia, 244; gracilis, 244
 Diandria, 54
 Dicksonia, 18
 Diceranaceae, 3
 Diceranella cerviculata, 1, 3; heteromalla, 3; secunda, 3
 Diceranum Bergeri, 4; Bonjeani, 4, var. juniperifolium, 4; elongatum, 1, 4; fuscescens, 4, var. congestum, 4, var. falcifolium, 4; groenlandicum, 1, 4; majus, 1, 4; scoparium, 4
 Didymodon rubellus, 5
 Digitaria, 49-52; The Grass Genus, 49
 Diphragma, sect. of Oxytropis, 147
 Distichium capillaceum, 3; Hagenii, 1, 3; inclinatum, 3
 Ditrichaceae, 3
 Draba, 77
 Drepanocladus fluitans, 10, var. falcatus, 10; revolvens, 10; uncinatus, 10, var. plumulosus, 10; vernicosus, 10
 Drew, K. M., Notes from the

Woods Hole Laboratory, 194;
 Trilliella intricata, 196
 Dumontia filiformis, 195

Eastern American Occurrence of
 Athyrium alpestre, 44; American
 Variety of Polystichum Braunii, 28

Eaton, R. J., Exceptional Dimensions in Corydalis sempervirens, 36; Exceptionally large Quercus ilicifolia, 19; Present Status of Magnolia virginiana in Massachusetts, 207

Ebony Spleenwort, 14
 Echinochloa, 26; colona, 51
 Edmondson, T. W., Solidago calcicola in Matane Co., Quebec, 157
 Ehrharta clandestina, 81
 Elaeagnus argentea, 122
 Eleocharis obtusa, 237; tenuis, 237
 Elodea canadensis, 39
 Empetrum, 77
 Encalypta rhabdocarpa, 4
 Encalyptaceae, 4
 Entodontaceae, 11
 Epidendrum, 54
 Epigeios, sect. of Calamagrostis, 202
 Epilobium, 137
 Equisetum, 79; arvense, 79, var. boreale, 79; hyemale, 79, var. affine, 80, f. intermedium, 80, f. pumilum, 80, var. intermedium, 80, var. Jesupi, 80, var. pumilum, 80; limosum, 79, f. natans, 79; litorale, 79; palustre, 79, var. americanum, 79, 80; scirpoides, 80; variegatum, 80, var. anceps, 80, var. Jesupi, 80

Eragrostis Damiensiana, 155-157; inconspicua, 156; peregrina, 155, 157, versus E. Damiensiana, 155; pilosa, 156, subsp. Damiensiana, 156, 157, var. Damiensiana, 156

Erica Tetralix, 204, 205
 Erigeron compositus, 122, var. multifidus, 122, var. trifidus, 122, var. typicus, 122; multifidus, 122; trifidus, 122

Erlanson, E. W., Ten New American Species and Varieties of Rosa, 109

Erucastrum Pollichii, 108
 Erythrocoma, 207; brevifolia, 207; ciliata, var. ornata, 207; flavula, 207

Eucephalus glaucescens, 228; glaucophyllus, 228; glaucus, 228; Gormanii, 228; vialis, 228

- Euphrasia*, 77
Eutriseta, sub-sect. *Trisetum*, 211, 212
 Evidence of the Hybrid Nature of *Betula Sandbergi*, 125
 Exceptional Dimensions in *Corydalis sempervirens*, 36
 Exceptionally large *Quercus ilicifolia*, 19

Farinosae, sect. of *Primula*, 60, 62, 63, 74, 86, 92, 96
 Fassett, N. C., *Bidens connata* and its Varieties in Wisconsin, 31; Notes from the Herbarium of the University of Wisconsin, II, 31, III, 205
 Fern, Beech, 16; Bladder, 18; Broad Beech, 16; Chain, 14; Christmas, 14; Hay-scented, 18; Lady, 15; Marsh, 16; Massachusetts, 16; New York, 16; Oak, 16; Ostrich, 18; Sensitive, 18; Shield, 16; Snake, 13; Wood, 16
 Fernald, M. L., American and eastern Asiatic *Beckmannia*, 24; American Representatives of *Asplenium Ruta-muraria*, 37; American Representative of *Calamagrostis Epigejos*, 202; Authors of *Sonchus arvensis*, var. *glabrescens*, 19; Eastern American Occurrence of *Athyrium alpestre*, 44; Eastern American Variety of *Polystichum Braunii*, 28; *Eragrostis peregrina* versus *E. Damiensiana*, 155; Genus *Oxytropis* in northeastern America, 137; *Geocaulon*, a New Genus of the *Santalaceae*, 21; New *Oxytropis* from the Gaspé Coast, A, 121; North American Species of *Anemone* § *Anemonanthea*, 180; *Panicum longifolium* in Massachusetts, 190; *Primula*, § *Farinosae* in America, 59, 85; *Roripa islandica* and *R. hispida*, 131; Some eastern American Forms of *Senecio*, 224; Varieties of *Galium boreale*, 106; Victorin's *Les Équisétinées* du Québec (review), 79
Festuceae, 242
 Fifth International Botanical Congress, Cambridge, 1930, 158
Fissidens osmundoides, 2
Fissidentaceae, 2
 Flora of Cape Cod, Additions to, 135; of Massachusetts, Reports on, I, 12; of Worcester Co., Mass., Additions to, 20
 Fogg, J. M., Jr., The Clandestine Form of *Leersia oryzoides*, 81
Fontinalaceae, 9
Fontinalis antipyretica, 9; *dalecarlica*, 9
Funaria hygrometrica, 5; *microstoma*, 5
Funariaceae, 5

Galium, 106; *boreale*, 106, The Varieties of, 106, var. *hyssopifolium*, 107, var. *intermedium*, 107, var. *scabrum*, 107, var. *typicum*, 107
 Gaspé Coast, A New *Oxytropis* from, 121
 Genus *Oxytropis* in northeastern America, 137; *Trisetum* in America, 209, 237
Geocaulon, 21, 23, A New Genus of the *Santalaceae*, 21; *lividum*, 23, 24
Geranium lobatum, 56; *triste*, 56
Gesneria, 54
Geum, 207; *ciliatum*, 206, 207; *triflorum*, 206, 207, f. *pallidum*, 207, var. *ciliatum*, 207, f. *flavulum*, 207, f. *ornatum*, 207
Glyceria leptostachya, 215; *microtheca*, 215; *montana*, 215
Gongrosira Debaryana, 194
Graphephorum, sub-sect. of *Trisetum*, 211, 240; *Brandegei*, 241; *Shearii*, 213
 Grass Genus *Digitaria*, 49
 Grigg, F. W., Reports on the Flora of Massachusetts, I, 12
Grimmia alpicola, 5, var. *latifolia*, 5; *apocarpa*, 5
Grimmiaceae, 5
 Griscom, L., *Carex Mitchelliana* and other rare Plants near Cohasset, Mass., 198
Gymnostomum rupestre, 4

Habenaria cristata, 190, 191; *viridis*, 47, var. *bracteata*, 47, var. *interjecta*, 47
 Hay-scented Fern, 18
Hedwigia albicans, 9
Hedwigiaceae, 9
Hepatica americana, 29; *nobilis*, 29
 Herbarium of the University of Wisconsin, Notes from, II, 31, III, 205
Hermodyctylus, 55
Heterolytrum, subgenus of *Trisetum*, 211, 212

- Heterotheca, 201, A New Variety from Texas, 200; subaxillaris, 201, var. **petiolaris**, 201
- Hippochaete, 79
- Hoff, A. C., *Trilliella intricata*, 196
- Homalocenchrus, 82
- Hybrid Nature of *Betula Sandbergi*, Evidence of, 125
- Hygrohypnum dilatatum, 10
- Hylocomiaceae, 11
- Hylocomium proliferum, 11
- Hymenostylium curvirostre, 5
- Hypericum canadense, 188-190, A Variety of, 188, var. **magninsulare**, 188-190
- Hypnaceae, 11
- Hypnum Bambergeri, 11; cupressiforme, 11; imponens, 11; pallescens, 11; Vaucheri, 11
- Ilex glabra, 198; opaca, 198
- Illinois Asters, Some, 77
- Iris, 55; Pseudacorus, 55
- Isolytrum, subgenus of *Trisetum*, 211, 244
- Isopterygium pulchellum, 11
- Itea virginica, 235
- Jabotapita, 56
- Jaceae, 59
- Josselyn Botanical Society, Announcement of Meeting, 124
- Juncus pervetus, 204, 205
- Kelsey, S. L., A New *Oxytropis* from the Gaspé Coast, 121
- Kempferia, 54, 55
- Kiaeria Blyttii, 4; Starkei, 4
- Kidder, N. T., *Chenopodium carinatum* and other Unusual Weeds, 158; *Lemna trisulca* at Isle au Haut, Maine, 121; Two Weeds New to Milton, Mass., 36
- Knowlton, C. H., Reports on the Flora of Massachusetts, I, 12; Tetramerism in *Trillium grandiflorum*, 105
- Koeleria, 241, 242
- Koeleriformia, subsect. of *Trisetum*, 211, 241
- Lachnanthes tinctoria, 35
- Lagopoda, 59
- Lagopus, 59
- Laminaria platymeris, 195
- Leersia hexandra, 82; monandra, 82; oryzoides, 81, 82, 84, The Clandestine Form of, 81, f. *clandestina*, 84, f. **inclusa**, 84, 85
- Lemna trisulca*, 121, at Isle au Haut, Maine, 121
- Leptobryum pyriforme, 6
- Leptodictyum riparium, 10
- Leskeaceae, 9
- Leskeella nervosa**, 9
- Leucobryaceae, 4
- Leucobryum glaucum, 4
- Leuconymphaea, 54, 57, 58
- Lewis, I. F., Notes from the Woods Hole Laboratory, 1928, 193
- Linnaean Species, Concerning the proper Identification of, especially those based on Material collected by Clayton, 232
- Lippia, 54
- Littlefield, E. W., An uncommon Association of Pines in Northern New York, 129
- Long, Bayard, Newfoundland Mosses collected by, in 1924-26, 1
- Lotoidea, 59
- Louis-Marie, The Genus *Trisetum* in America, 209, 237
- Lupinaster, 59
- Lupulina, 59
- Lupulinum, 59
- Luzula pilosa, 29; saltuensis, 29
- Lycopodium annotinum, 198; clavatum, 135, 198; lucidulum, 136, 198
- Mackenzie, K. K., Concerning the proper Identification of Linnaean Species, especially those based on Material collected by Clayton, 232; The Grass Genus *Digitaria*, 49; *Solidago flexicaulis* and *Solidago latifolia*, 223
- Madia sativa, 36
- Magnolia virginiana, 207, Present Status of in Massachusetts, 207
- Maidenhair, 13
- Maine, *Lemna trisulca* at Isle au Haut, 121
- Malaxis unifolia, 199
- Marsh Fern, 16
- Maryland, *Aster ericoides*, var. *platyphyllus* in, 106
- Massachusetts, Additions to the Flora of Worcester Co., 20; *Carex Mitchelliana* and other rare Plants near Cohasset, 198; Fern, 16; *Panicum longifolium* in, 190; Present Status of *Magnolia virginiana* in, 207; Reports on the Flora of, I, 12; Two Weeds New to Milton, 36; *Utricularia subulata* in Plymouth, 35

- Meesia trichodes*, 8, var. *angustifolia*, 8
Meesiaceae, 8
Melica triflora, 238
Meliloti, 59
Melilotus, 59
Merismopedia elegans, 195; *tenuissima*, 195
Mesembryanthemum loreum, 56; *scabrum*, 55; *tortuosum*, 56
Microspora, 194
Minutissimae, sect. of *Primula*, 62
Mniaceae, 8
Mnium *hornum*, 2, 8; *punctatum*, var. *elatum*, 8; *rostratum*, 8; *subglobosum*, 8
 Missouri, a *Ranunculus* from, 200
 Mosses collected in Newfoundland by Mr. Bayard Long in 1924-26, 1
Myurella julacea, 9

Najas guadalupensis, 136
Narcissus Jonquilla, 69
Nasturtium hispidum, 133; *palustre*, 132, 133, var. *hispidum*, 133; *terrestre*, 132, 133
Nelumbo, 54, 56-58
Nemorosa quinquefolia, 186
Nenuphar album, 57; *luteum*, 57
Nephrodium marginale, 17, f. *Davenportii*, 17; *Thelypteris*, f. *Pufferae*, 16
Nestronia, 22, 23; *umbellula*, 22
 New England, A Note on the Genus *Paspalum* in, 133
 Newfoundland, Mosses collected by Mr. Bayard Long in, 1924-26, 1
 New Genus of the *Santalaceae*, A, 21
 New Hampshire, Plants New to Coos County, 104
 New *Oxytropis* from the Gaspé Coast, A, 121
 New York Fern, 16
 New York, Uncommon Association of Pines in Northern, 129
 New Variety of *Aster multiflorus*, 77
 North American Species of *Anemone* § *Anemonanthea*, 180
 Note on the Genus *Paspalum* in New England, 133
 Notes from the Herbarium of the University of Wisconsin, II, 31, III, 205; from the Woods Hole Laboratory, 1928, 193; on *Aster*, 226; on certain Species of *Panicum* occurring in or near Rhode Island, 229

Nymphaea, 53, 54, 56-58, Standard Species of, 53; *alba*, 54, 56-58; *lutea*, 53, 54, 56, 57

 Oak Fern, 16
Ochna Jabotapita, 56
Oedogonium, 193; *Reinschii*, 193
Oncophorus virens, 4; *Wahlenbergii*, 3
Onoclea, 18; *sensibilis*, 18, f. *obtusilobata*, 18; *Struthiopteris*, 18
Oreoweisia serrulata, 3
Orthodieranum flagellare, 4; *montanum*, 4
Orthotrichaceae, 9
Orthotrichum anomalum, 9; *Blyttii*, 9
Oryza, 81, 82; *clandestina*, 82, 83, f. *inclusa*, 83, 84, f. *patens*, 83; *oryzoides*, 82
 Ostrich Fern, 18
Oxalis europaea, f. *villicaulis*, 158
Oxytropis, 121-123, 137, 138, 145, 150, 151, from the Gaspé Coast, A New, 121, in northeastern America, The Genus, 137; *arctica*, 139, 143-145, 150, 152, 155, δ *inflata*, 154, 155, β *minor*, 125, α *subumbellata*, 144, var. *uniflora*, 152; *arctobia*, 139, 145, 152, 153; *Bellii*, 139, 145, 150, 152; *Bellii*, 150, 151; *borealis*, 144; *campestris*, 141-144, 146-150, 155, var. *caerulea*, 142, 146, 148, 149, var. *johannensis*, 146, var. *melanocephala*, 142-144, 155, var. *sordida*, 143, 152; ***coronaminis***, 151, 152; *deflexa*, 140, 141, β *sericea*, 140; *foliolosa*, 138-140; *foliosa*, 140; ***gaspensis***, 123, 138, 141; *gracilis*, 123, 124, 141; *Hallii*, 154, 155; ***hudsonica***, 138, 142, 145; ***johannensis***, 139, 145-147, 149, 150; *Lamberti*, 146, 149, var. *sericea*, 146; *lapponica*, 140, 141; *leucantha*, 142-144; *Maydelliana*, 139, 142-145; *nigrescens*, 153, var. *arctobia*, 152; *podocarpa*, 139, 150, 154, 155; ***pygmaea***, 153; ***retrorsa***, 140, 141, var. ***sericea***, 140, 141; *Roldi*, 152; *sordida*, 148, 149, 152; ***terrae-novae***, 139, 147, 149; *uralensis*, 146, γ *minor*, 148, var. *pumila*, 144; *viscida*, 122-124, 141; *viscidula*, 141

Paniceae, 215
Panicum, 229, Notes on certain

- Species of, occurring in or near Rhode Island, 229; Addisonii, 230; albemarleense, 230; amarum, 230; annulum, 230; auburne, 230; barbipulvinatum, 230; barbula-tum, 230; Bicknellii, 230; Boscii, 230, var. molle, 230; brizoides, 26; calliphyl-lum, 230; capillare, 230, var. occidentale, 230; Clutei, 230; colonum, 51; columbianum, 230, var. thinium, 230; Common-sianum, 230; commutatum, 230; depauperatum, var. psilophyl-lum, 231; dichotomiflorum, 231, var. puritanorum, 231; dichoto-mum, 231; dissectum, 51; lanu-ginosum, 231; linearifolium, 231, var. Wernerii, 231; longifolium, 190, 191, 231, in Massachusetts, 190, var. tusketense, 191; lucidum, 230; mattamuskeetense, 231; miliaceum, 231; minutulum, 35, 230; oligosanthos, 231; philadel-phicum, 231; pseudopubes-cens, 230; scoparioides, 231; sco-parium, 231; stipitatum, 230; subvillosum, 231; Syzigachne, 26, 27; villosissimum, 232; Wrighti-anum, 230; xanthophyllum, 230
- Paraleucobryum longifolium*, 3
- Paspalum*, 52, 133, A Note on the Genus in New England, 133; ciliatifolium, 134; circulare, 134, 135; Muhlenbergii, 134; pleni-pilum, 134, 135; psammophilum, 135; pubescens, 134, 135; seta-ceum, 133-135; virgatum, 51, 52
- Pease, A. S., Plants New to Coos County, New Hampshire, 104; Reports on the Flora of Massa-chusetts, I, 12
- Pellaea*, 14; atropurpurea, 14
- Perezia aletes*, 20
- Phalaris erucaeformis*, 24, 25, 51; oryzoides, 81, 82
- Phegopteris*, 16, 48; alpestris, 44, 48, 49; *Dryopteris*, 16; hexagon-optera, 16; polypodioides, 16
- Phillyrea angustifolia*, 55, 56; media, 55
- Philonotis fontana*, 9
- Phragmites*, 202
- Phyllophora membranifolia*, 196
- Pine, Jack, 129; Pitch, 129-131; Red, 129, 130; White, 129
- Pines, 129, An uncommon Associa-tion of, in Northern New York, 129
- Pinus Banksiana*, 129, 130; *resinosa*, 129; *rigida*, 129, 130; *Strobus*, 129
- Plagiopus Oederi*, 8
- Plagiotheciaceae, 11
- Plagiothecium denticulatum*, 11; *silvaticum*, 11; *striatellum*, 11
- Plantago juncoideis*, 77
- Plants New to Coos County, New Hampshire, 104
- Pleurozium Schreberi*, 11
- Poa tenuifolia*, 215
- Pogonatum capillare*, 2, 12; *urni-gerum*, 12
- Polemonium micranthum*, 36
- Polygonum glaucum*, 204; *mariti-mum*, 204
- Polypodiaceae, 13
- Polypodium*, 13, 48; *rhaeticum*, 48; *virginianum*, 13, 136, f. *acumi-natum*, 13, f. *bipinnatifidum*, 13, f. *deltoideum*, 13; *vulgare*, 13, var. *attenuatum*, 13, var. *auri-tum*, 13, var. *cambricum*, 13
- Polypody, 13
- Polystichum aculeatum*, var. *Braun-nii*, 30; *acrostichoides*, 16, 136, f. *crispum*, 16, f. *incisum*, 16, var. *Schweinitzii*, 16; *alaskense*, 28; *Braunii*, 16, 28-30, The east-ern American Variety of, 28, var. *japonicum*, 28, var. *kamtschati-cum*, 28, var. **Purshii**, 30; *hale-akalense*, 28; *Hillebrandii*, 28
- Polytrichaceae, 12
- Polytrichum*, 204; *alpinum*, 12, var. *silvaticum*, 12; *commune*, var. *perigoniale*, 12; *gracile*, 12; *juniperinum*, 12; *ohioense*, 12; *piliferum*, 12; *strictum*, 12
- Poole, J. P., *Oedogonium Rein-schii*, 193
- Potentilla*, 104; *argentea*, 107; *intermedia*, 107, in the Boston District, 107; *pulcherrima*, 105
- Pottia Heimii*, 5
- Pottiaceae, 4
- Present Status of *Magnolia vir-giniana* in Massachusetts, 207
- Primula*, 59; *ajanensis*, 65, 88, 102; *americana*, 73, 86; *arctica*, 63; *borealis*, 65, 67, 88, 94-99, 102, var. *Loczii*, 95, 96, var. *parvi-folia*, 94, 95; *Chamissonis*, 94; *dealbata*, 73, 74; *decipiens*, 60, 61, 65, 74-77, 102; *egaliccensis*, 101; *egalikensis*, 65-68, 96-102, f. **violacea**, 98, 99, 102; *egaliccensis*, 97; *eximia*, 63; *farinosa*, 60-62, 66-77, 85-87, 99, β *ameri-*

- cana, 60, 86, 87, β stricta, 66, subsp. eufarinosa, var. genuina, 61, 69, 73, subsp. groenlandica, 99, subsp. mistassinica, 89, var. americana, 61, 69, 71, var. nudata, 72, var. genuina, 69, 73, var. groenlandica, 66-68, 98, var. incana, 73, var. macropoda, 61, 62, 66, 69, 70, 73, var. magellanica, 74-76, var. mistassinica, 66, 89; § Farinosae, 59, 60, 62-64, 74, 77, 85, in America, 59, 85; finmarchica, 100; glabrescens, 66; groenlandica, 99; Hornemania, 66, 89; incana, 60-62, 64, 71, 72, 74, 76, 77, 87, 102; **intercedens**, 65, 77, 86, 87, 102; integrifolia, 101, 102; intermedia, 101; **laurentiana**, 64, 68, 70-72, 77, 87, 102, f. **chlorophylla**, 72, 102; Loczii, 94; Loczyi, 95-97; Maccalliana, 89, 92, 93; Macounii, 63; macropoda, 69, 70; magellanica, 61, 74-76; mistassinica, 60-62, 65-67, 71, 72, 87-95, 98, 99, 102, f. leucantha, 60, 88, 89, 91, 102, var. **novoboracensis**, 88, 89, 91-93, 102, var. **typica**, 89; nivalis, 63, γ pumila, 63; norvegica, 100, 101; parvifolia, 65, 93, 94, 102; pumila, 63, var. arctica, 63, var. Ledebouriana, 63; pusilla, 60, 69, 86, 89; rotundifolia, 101, 102; Rusbyi, 86; scotica, 67, 69; sibirica, 65-67, 97, 98, 100-102, γ minor, 97, var. arctica, 99, var. borealis, 95, var. β integrifolia, 101, 102, var. kashmiriana, 102, var. mistassinica, 89, var. rotundifolia, 102; specuicola, 60, 65, 85, 86, 102; stricta, 61, 64, 66-68, 72, 74, 77, 87, 88, 95, 98, 99, 102, var. groenlandica, 66-68, 98, 99, 101, var. jacutensis, 66, 67; tenuis, 94; Tschuktschorum, 63, var. **arctica**, 63, var. **pumila**, 63
 Primulaceae, 94
 Priva, 54
 Prunus, 235; caroliniana, 235; nana, 235; serotina, 235; virginiana, 234-236
 Pseudathyrium, 48; alpestre, 48
 Pteretis, 18; nodulosa, 18
 Pteridium, 13; latiusculum, 13, var. pseudocaudatum, 13
 Pteris aquilina, 13, var. pseudocaudata, 13
 Ptilium crista-castrensis, 11
 Puccinellia, 77
 Quebec, Solidago calcicola in Matane County, 157
 Quercus coccinea, 235; ilicifolia, 19, An exceptionally large, 19; rubra, 235
 Radicula, 132; hispida, 133; palustris, 132, 133, var. hispida, 133
 Ranunculus, 200, A New, from Missouri, 200; abortivus, 200; alleghaniensis, 201; Harveyi, 200, var. **pilosus**, 200
 Reports on the Flora of Massachusetts, I, 12
 Rhacomitrium canescens, 5; fasciculare, 5; heterostichum, var. gracilescens, 5; hypnoides, 5
 Rhapontica, 59
 Rhaponticoides, 59
 Rhaponticum, 59
 Rhode Island, Notes on certain Species of Panicum occurring in or near, 229
 Rhus, 55
 Rhytidiaceae, 11
 Rhytidiadelphus loreus, 11; squarrosus, 11; triquetrus, 11
 Rhytidium rugosum, 11
 Robinia Pseudacacia, 203
 Rock Brake, 14
 Roripa, 132; hispida, 131, 133, var. glabrata, 133; islandica, 131, 132; palustris, 131-133, var. hispida, 131-133; sylvestris, 131
 Rosa, 109, Ten New American Species and Varieties of, 109; arkansana, 116; blanda, 110-112, 114, 117, var. **Hermannii**, 113, var. nuda, 110, var. subgeminata, 110; Bushii, 110, 117; carolina, 110, 111, 113, 117, 119, 121, \times humilis, 113, \times palustris, 113, var. aculeata, 110, var. **litoralis**, 119, var. **sabulosa**, 118, var. sepalorelevata, 110; conjuncta, 110; **Deamii**, 120; **Housei**, 111, 117; **Lyoni**, 121; **michiganensis**, 111; obovata, 121; palustriformis, 110; palustris, 110-114, var. **aculeata**, 110, var. **sepalorelevata**, 110; polyanthema, 110; pratincola, 116; **ratonensis**, 115, 116; **relicta**, 116, 117; rudiusscula, 110; **Schuettea**, 112; suffulta, 110, 111, 115-118, var. **valida**, 114, 116; Woodsii, 116

- Rosendahl, C. O., Evidence of the Hybrid Nature of *Betula Sandbergii*, 125
Rumex maritimus, var. *fueginus*, 108
Rynchospora inundata, 136
Salix, 137
Salsola Kali, var. *tenuifolia*, 158
Sanford, S. N. F., Reports on the Flora of Massachusetts, I, 12
Santalaceae, 21, 23; *Geocaulon*, a New Genus of, 21
Satyrion, 54
Saxifraga, 77
Scirpus capitatus, 237
Scorpidium scorpioides, 10
Seiropora Griffithsiana, 195
Seligeria Donniana, 3
Seligeraceae, 3
Sematophyllaceae, 11
Senecio, 224, Some eastern American Forms of, 224; *aureus*, 225; *Burkei*, 225; *elongatus*, 225; *Fernaldii*, 225; *hyperborealis*, var. *columbiensis*, 225; *indecus*, f. *Burkei*, 225; *obovatus*, f. *elongatus*, 225; *pauciflorus fallax*, 225, f. *fallax*, 225, var. *fallax*, 225; *pauperulus*, 226, f. *verecundus*, 225, var. *Balsamitae*, f. *inchoatus*, 226; *pseudoreus*, 225, f. *ecoronatus*, 225; *resedifolius*, f. *columbiensis*, 225, var. *columbiensis*, 225; *Robbinsii*, f. *invenustus*, 225
Sensitive Fern, 18
Sericocarpus asteroides, 136, f. *roseus*, 136
Sherardia, 54
Shield Fern, 16
Sieversia, 207; *ciliata*, 207
Silvery Spleenwort, 15
Sisymbrium amphibium, 132, α *palustre*, 132, δ *terrestre*, 132; *hispidium*, 133; *islandicum*, 132; *palustre*, 132; *terrestre*, 132
Sisyrinchium, 55
Smith, L. B., Reports on the Flora of Massachusetts, I, 12; *Utricularia subulata* in Plymouth, Mass., 35
Snake Fern, 13
Solidago caesia, 223; *calicicola*, 105, 157, in Matane County, Quebec, 157; *flexicaulis*, 223, 224; *lateriflora*, 172, 176, 224; *latifolia*, 223, 224; *macrophylla*, 105, 157; *menalis*, 157
Some Eastern American Forms of *Senecio*, 224; Illinois Asters and a New Variety of *A. multiflorus*, 77
Sonchus arvensis, 19, var. *glabrescens*, 19, Authors of, 19, var. *laevipes*, 19
Spermothamnion roseolum, 196; *Turneri*, 197
Sphagnaceae, 2
Sphagnum capillaceum, 2; *imbricatum*, 2; *Lindbergii*, 2; *palustre*, 2; *pulchrum*, 2; *Pylaeii*, 2; *subsecundum*, 2; *tenellum*, 2; *Warnstorffii*, 2
Sphenopholidea, sub-sect. of *Trisetum*, 211, 240
Spiesia arctica, 144; *Belli*, 150; *campestris*, 146, 148; *inflata*, 154; *podocarpa*, 154
Splachnaceae, 6
Splachnum ampullaceum, 6; *pedunculatum*, 6
Spleenwort, 14, 15; *Ebony*, 14; *Silvery*, 15
Sprague, T. A., The Standard Species of *Nymphaea* L., 53
Standard Species of *Nymphaea*, 53
Stegonia latifolia, 1, 5
Stoebae, 59
Stachytarpheta, 54
Svenson, H. K., Additions to the Flora of Cape Cod, 135; *Carex Mitchelliana* and other rare Plants near Cohasset, Mass., 198
Syntherisma sanguinale, 50, 51
Talinum rugospermum, 205, 206; *teretifolium*, 205, 206
Tamonea, 54
Taraxacum, 77, 137
Taylor, W. R., Notes from the Woods Hole Laboratory, 1928, 193
Tayloria splachnoides, 6
Ten New American Species and Varieties of *Rosa*, 109
Tetramerism in *Trillium grandiflorum*, 105
Tetraplodon angustatus, 6; *bryoides*, 6, var. *cavifolius*, 6, var. *paradoxus*, 6
Texas, A *Heterotheca* from, 200
Theliaceae, 9
Thelypteris Boottii, 17, 18; *cristata*, 17, var. *Clintoniana*, 17; *Dryopteris*, 16; *Gouldiana*, 17; *hexagonoptera*, 16; *marginalis*, 17, f. *Davenportii*, 17, f. *elegans*,

- 17; noveboracensis, 16; palustris, 16, f. **Pufferae**, 16; Phegopteris, 16, 136; simulata, 16; spinulosa, 17, var. americana, 17, 29, var. concordiana, 17; var. dilatata, 29, var. frutuosa, 17, var. intermedia, 17
- Thuidiaceae, 9
- Thuidium delicatulum, 9
- Tomenthypnum nitens, 10
- Tortella tortuosa, 5
- Tortula norvegica, 5; ruralis, 5
- Toxicodendron, 55
- Trailiella intricata, 196, 197
- Tribonema, 194
- Trifolium, 59
- Trifolium, 59
- Trillium grandifolium, 105, Tetramerism in, 105
- Triphyllodes, 59
- Tripsacum dactyloides, 51
- Trisetum, 209, 210, 237, 241, 242, The Genus in America, 209, 237; airoides, 238; alaskanum, 221, 239; altijugum, 241; **andicola**, 244; andinum, 238, 239; antarcticum, 240; araeanthum, 242; argenteum, 213; bambusiforme, 212; barbatum, 243, var. majus, 243; barbinode, 240, var. **hirtiflorum**, 240; biflorum, 239; **Bongardii**, 220; brachyatherum, 242; Brandegei, 241; brasiliense, 242; Brittonii, 239; Buchtienii, 238; californicum, 240; canescens, 214-216, 218, f. **tonsum**, 216, f. **velutinum**, 216; caudulatum, 240; cernuum, 213-217, 219, var. **luxurians**, 213, f. **pubescens**, 213, var. **Sandbergii**, 214; chromostachyum, 240; chiloense, 240; Congdoni, 219, 239, 240; depauperatum, 242; deyeuxioides, 244; **disjunctum**, 218; elatum, 215-217; erectum, 240; evolutum, 245; filifolium, 217, var. aristatum, 217; flavescens, 212, 216; floribundum, 243, var. **Weberbaueri**, 244; Fournieranum, 239; fraudulentum, 212; groenlandicum, 238; Hallii, 240; heteronymum, 217; hirsutum, 239; hirtiflorum, 240; hirtum, 217; humile, 244; interruptum, 218, 240, var. **californicum**, 240; irazuense, 212; Jungensii, 243; labradoricum, 238; lasiolepis, 238; laxiflorum, 242; laxum, 242; litorale, 243; longiglume, 244; Macbridei, 245; malacophyllum, 212, 218; melicoides, 240; micratherum, 242; **mollifolium**, 218; montanum, 212, var. **pilosum**, 212, var. **Shearii**, 213; monticola, 240; muticum, 241; nemorosum, 242; nivolum, 239; ochrostachyum, 240; **oreophilum**, 221, 223, var. **Johnstonii**, 237; Palmeri, 242; paniculatum, 212; paradoxum, 240; pennsylvanicum, 240; Preslii, 238, var. Buchtienii, 238, var. **lasiantha**, 238; Pringlei, 241; **projectum**, 217; Rosei, 223, 239, f. **tenerum**, 239, var. **tenerum**, 239; Sandbergii, 214; sesquiflorum, 219, 220, 239, 240; Shearii, 213; spicatum, 219, 223, 238, 239, var. **alaskanum**, 239, var. **andinum**, 239, var. **Brittonii**, 239, var. **dianthemum**, 239, var. **fuegianum**, 239, var. **hirsutum**, 239, var. **laxius**, 239, var. Maidenii, 239, var. majus, 239, var. molle, 239, var. **nivolum**, 239, var. phleoides, 239, var. pilosiglume, 239, var. **villosissimum**, 239; splendidulum, 212; subspicatum, 238, var. breviglume, 238, var. compactum, 239, var. fuegianum, 239, var. glabrifolium, 238, var. laxius, 239, var. muticum, 241, var. villosissimum, 239; tolucense, 238; **Trinii**, 243, var. **effusum**, 243, var. **litorale**, 243, var. **majus**, 243, var. **manicatum**, 243, var. **micrantherum**, 243, var. **pallidiflorum**, 243, var. **strictum**, 243; variabile, 240, var. **chiloense**, 240, var. **flavescens**, 240, var. **intonsum**, 240, var. **Vidali**, 240, var. **virescens**, 240; **Vidali**, 240; viride, 244; Virletii, 212; Weberbaueri, 244; **Williamsii**, 221; Wolfii, 240, var. **Brandegei**, 241, f. **muticum**, 241, var. **muticum**, 241
- Two New Varieties of Early Spring Plants, a Ranunculus from Missouri and a Heterotheca from Texas, 200; Weeds New to Milton, Mass., 36
- Ulotia americana, 9; crispula, 9; Ludwigii, 2, 9; phyllantha, 2, 9
- Uncommon Association of Pines in Northern New York, 129
- University of Wisconsin, Notes

- from the Herbarium of, II, 31;
III, 205
Utricularia, 35; geminiscapa, 121;
subulata, 35, in Plymouth, Mass.,
35, f. cleistogama, 35
Vaccinium Oxycoccus, 71; uliginosum, 71
Varieties of Galium boreale, 106;
of Ranunculus and Heterotheca,
New, 200; of Rosa, New American, 109
Variety of Hypericum canadense,
188; of Polystichum Braunii, The
eastern American, 21
Verbena, 54; officinalis, 54
Vernix, 55
Vesicaria, 59
Victorin's Equisétinées du Québec
(review), 79
Walking Leaf, 15
Weatherby, C. A., Additions to the
Flora of Worcester Co., Mass
(notice), 20; Note on the Genus
Paspalum in New England, 133;
Variety of Hypericum canadense,
188
Webera commutata, 6; cruda, 6;
nutans, 6; prolifera, 6
Weeds, Chenopodium carinatum
and other unusual, 158; Two New
to Milton, Mass., 36
Wiegand, K. M., Aster lateriflorus
and some of its Relatives, 161
Wisconsin, Bidens connata and its
Varieties in, 31
Wood Fern, 16
Woods Hole Laboratory, Notes
from, 1928, 193
Woodsia, 18; alpina, 18; ilvensis,
18; obtusa, 18
Woodwardia, 14; areolata, 14; virginica, 14
Xiphium, 55
Xylosteum villosum, 90
Xyris Smalliana, 35

DUPLICATE BOOKS FOR SALE

- Gordon, George.** The pinetum. A Synopsis of all the Coniferous Plants at present known with descriptions, history and synonyms. 1858. 8vo., 353 pp..... **\$2.00**
 Same. Supplement. 1862. 8vo. 119 pp.... **\$1.00**
- Loudon, John C.** Encyclopaedia of plants. New Edition. Edited by Mrs. Loudon. 1855. 8vo., 1574 pp. With over 20,000 figures. Half calf **\$3.50**
- Meehan, Thomas.** The native flowers and ferns of the United States. 2 series in 4 vols. With 192 colored plates. Boston and Philadelphia. 1878-80. Large 8vo. Cloth..... **\$14.00**
- Sullivant, W. S.** Icones Muscorum, 8vo., illustrated by numerous copper plates of high excellence (unbound and in sheets). Here offered at a greatly reduced price..... **\$6.00**
 Supplement to the preceding (also in sheets). **\$5.00**
 Icones and Supplement together..... **\$10.00**

Address Librarian, GRAY HERBARIUM of HARVARD UNIVERSITY, Cambridge, Mass

Chinese Economic Trees by Professor Woon Young Chun. 318 pp., 100 outline illustrations. Shanghai, 1921. \$3.15 postpaid.

Catalogue of Trees and Shrubs of China by Professor H. H. Chung. 271 pp., Shanghai, 1924, \$1.10 postpaid.

Apply to J. G. JACK, East Walpole, Massachusetts.

FIELD AND HERBARIUM EQUIPMENT

HAND LENSES	COLLECTING CASES	PLANT PRESSES
FERN TROWELS	MOUNTING PAPERS	GENUS COVERS
FIELD PICKS	HERBARIUM CASES	FELT DRIERS

Write for free Catalog R-91

CAMBRIDGE BOTANICAL SUPPLY CO., Waverley, Mass.